

FILTER GAMES AND COMBINATORIAL PROPERTIES OF WINNING STRATEGIES

CLAUDE LAFLAMME

ABSTRACT. We characterize winning strategies in various infinite games involving filters on the natural numbers in terms of combinatorics or structural properties of the given filter. These generalize several ultrafilter games of Galvin.

1. INTRODUCTION

We look at various infinite games between two players **I** and **II** involving filters on the natural numbers in which **I** either plays cofinite sets, members of $\mathcal{F}$ or $\mathcal{F}^+$ and player **II** responds with an element or a finite subset of **I**'s move, depending on the game. In each version, the outcome depends on the set produced by player **II**, whether it belongs to the given filter $\mathcal{F}$, $\mathcal{F}^+$ or even $\mathcal{F}^*$, the dual ideal.

In each game considered, we will characterize winning strategies of either player in terms of combinatorics of the given filter $\mathcal{F}$; these combinatorics turn out to be generalizations of the classical notions of P-points, Q-points and selectivity for ultrafilters. In the case of ultrafilters, $\mathcal{F} = \mathcal{F}^+$ and most of our games to ones already studied by Galvin in unpublished manuscripts [5]; the various generalized combinatorics enjoyed by the filters become equivalent.

Several characterizations of Ramsey ultrafilters and P-points were known from works of Booth ([4]) and Kunen ([8]), and some generalizations of these combinatorics to filters were already made by Grigorieff in [6] where for example the notion of P-filter is characterized in terms of branches through certain trees; we shall see that this is very much in the spirit of winning strategies for certain games.

Other variations of these games for ultrafilters can be found in Chapter VI of Shelah's book [11], and two of the games below have been analyzed by Bartoszynski and Scheepers [2].

We wish to thank Chris Leary for helpful suggestions and corrections regarding the present paper.

Our terminology is standard but we review the main concepts and notation. The natural numbers will be denoted by ω , $\wp(\omega)$ denotes the collection of all its subsets. Given $X \in \wp(\omega)$, we write $[X]^\omega$ and $[X]^{<\omega}$ to denote the infinite or finite subsets of X respectively. We use the well known 'almost inclusion' ordering between members of $[\omega]^\omega$, i.e. $X \subseteq^* Y$ if $X \setminus Y$ is finite. We identify $\wp(\omega)$ with ${}^\omega 2$ via characteristic functions. The space ${}^\omega 2$ is further equipped with the product topology of the discrete space $\{0, 1\}$. A basic neighbourhood is then given by sets of the form

Date: April 12, 1994.

1991 *Mathematics Subject Classification.* Primary 04A20; Secondary 03E05, 03E15, 03E35.

This research was partially supported by NSERC of Canada.

$$\mathcal{O}_s = \{f \in {}^\omega 2 : s \subseteq f\}$$

where $s \in {}^{<\omega} 2$, the collection of finite binary sequences. The terms “nowhere dense”, “meager”, “Baire property” all refer to this topology. Concatenation of elements $\bar{s}, \bar{t} \in {}^{<\omega} \omega$ will be written $\bar{s} \wedge \bar{t}$.

A filter is a collection of subsets of ω closed under finite intersections, supersets and containing all cofinite sets; it is called proper if it contains only infinite sets. For a filter $\mathcal{F}$, $\mathcal{F}^+$ denotes the collection of all sets X such that $\langle \mathcal{F}, X \rangle$ is a proper filter; it is useful to notice that $X \in \mathcal{F}^+$ if and only if $X^c \notin \mathcal{F}$. $(\mathcal{F}^+)^c = \wp(\omega) \setminus \mathcal{F}^+$, the collection of sets incompatible with $\mathcal{F}$ is the dual ideal and is usually denoted by $\mathcal{F}^*$. The *Fréchet* filter is the collection of cofinite sets, denoted by $\mathfrak{Fr}$.

The families $\mathcal{F}$ and $\mathcal{F}^+$ are dual in a different sense; this means that a set X containing an element of each member of $\mathcal{F}$ (resp. $\mathcal{F}^+$) must belong to $\mathcal{F}^+$ (resp. $\mathcal{F}$). In particular $\mathfrak{Fr}$ and $[\omega]^\omega$ are dual. From more general work of Aczel ([1]) and Blass ([3]), there is a duality between games in which a player chooses $X_k \in \mathcal{F}$ while the other player responds with $n_k \in X_k$, and games in which a player chooses $Y_k \in \mathcal{F}^+$ while the other player responds with $n_k \in Y_k$. The point is that the statements

$$(\forall X \in \mathcal{F})(\exists n \in X)\phi(n) \text{ and } (\exists Y \in \mathcal{F}^+)(\forall n \in Y)\phi(n)$$

are equivalent.

The following important result characterizes meager filters in terms of combinatorial properties.

Proposition 1.1. (Talagrand ([12])) *The following are equivalent for a filter $\mathcal{F}$:*

1. $\mathcal{F}$ has the Baire property.
2. $\mathcal{F}$ is meager.
3. There is a sequence $n_0 < n_1 < \dots$ such that

$$(\forall X \in \mathcal{F})(\forall^\infty k) X \cap [n_k, n_{k+1}) \neq \emptyset.$$

Combinatorial properties of filters have played an important role in applications of Set Theory, and the classical notions of a filter being meager, a P-filter or selective have been around a long time. These concepts will be generalized below in terms of trees and other structural properties; these combinatorial ideas have their roots in Ramsey theory and P-points and selective ultrafilters (sometimes called ‘Ramsey’) have characterizations in term of these trees; this can be found in the papers by Booth [4] and Grigorieff [6].

We call a tree $\mathcal{T} \subseteq {}^{<\omega} \omega$ an $\mathcal{X}$ -tree for some $\mathcal{X} \subseteq [\omega]^\omega$ ($\mathcal{X}$ will usually be a filter $\mathcal{F}$ or $\mathcal{F}^+$), if for each $\bar{s} \in \mathcal{T}$, there is an $X_{\bar{s}} \in \mathcal{X}$ such that $\bar{s} \wedge n \in \mathcal{T}$ for all $n \in X_{\bar{s}}$. Similarly we call a tree $\mathcal{T} \subseteq {}^{<\omega} ([\omega]^{<\omega})$ an $\mathcal{X}$ -tree of finite sets for some $\mathcal{X} \subseteq [\omega]^\omega$, if for each $\bar{s} \in \mathcal{T}$, there is an $X_{\bar{s}} \in \mathcal{X}$ such that $\bar{s} \wedge a \in \mathcal{T}$ for each $a \in [X_{\bar{s}}]^{<\omega}$. A branch of such a tree is thus an infinite sequence of finite sets and we will be interested in the union of such a branch, an infinite subset of ω .

Here are a few more combinatorial properties of filters that we will consider.

Definition 1.2. *Let $\mathcal{F}$ be a filter on ω .*

1. $\mathcal{F}$ is called a *Q-filter* if for any partition of ω into finite sets $\langle s_k : k \in \omega \rangle$, there is an $X \in \mathcal{F}$ such that $|X \cap s_k| \leq 1$ for all k .

2. $\mathcal{F}$ is called a weak Q -filter if for any partition of ω into finite sets $\langle s_k : k \in \omega \rangle$, there is an $X \in \mathcal{F}^+$ such that $|X \cap s_k| \leq 1$ for all k .
3. $\mathcal{F}$ is called diagonalizable if there is an $X \in [\omega]^\omega$ such that $X \subseteq^* Y$ for all $Y \in \mathcal{F}$.
4. $\mathcal{F}$ is called ω -diagonalizable if there are $\langle X_n \in [\omega]^\omega : n \in \omega \rangle$ such that for each $Y \in \mathcal{F}$, there is an n such that $X_n \subseteq^* Y$.
5. $\mathcal{F}$ is called ω -+-diagonalizable if there are $\langle X_n \in \mathcal{F}^+ : n \in \omega \rangle$ such that for each $Y \in \mathcal{F}$, there is an n such that $X_n \subseteq^* Y$.
6. A set $X \subseteq [\omega]^{<\omega}$ is called $\mathcal{Z}$ -universal ($\mathcal{Z}$ will be $\mathcal{F}$ or $\mathcal{F}^+$) if for each $Y \in \mathcal{Z}$, there is an $x \in X \cap [Y]^{<\omega}$. $\mathcal{F}$ is called ω -diagonalizable by $\mathcal{Z}$ -universal sets if there are $\mathcal{Z}$ -universal sets $\langle X_n : n \in \omega \rangle$ such that for all $Y \in \mathcal{F}$, there is an n such that $x \cap Y \neq \emptyset$ for all but finitely many $x \in X_n$.
7. $\mathcal{F}$ is a P -filter if given any sequence $\langle X_n : n \in \omega \rangle \subseteq \mathcal{F}$, there is an $X \in \mathcal{F}$ such that $X \subseteq^* X_n$ for each n .
8. $\mathcal{F}$ is a weak P -filter if given any sequence $\langle X_n : n \in \omega \rangle \subseteq \mathcal{F}$, there is an $X \in \mathcal{F}^+$ such that $X \subseteq^* X_n$ for each n . Equivalently, every $\mathcal{F}$ -tree of finite sets has a branch whose union is in $\mathcal{F}^+$.
9. $\mathcal{F}$ is a P^+ -filter if every $\mathcal{F}^+$ -tree of finite sets has a branch whose union is in $\mathcal{F}^+$.
10. $\mathcal{F}$ is Ramsey if any $\mathcal{F}$ -tree has a branch in $\mathcal{F}$; equivalently, $\mathcal{F}$ is both a Q -filter and a P -filter.
11. $\mathcal{F}$ is weakly Ramsey if any $\mathcal{F}$ -tree has a branch in $\mathcal{F}^+$.
12. $\mathcal{F}$ is +-Ramsey if every $\mathcal{F}^+$ -tree has a branch in $\mathcal{F}^+$.
13. $\mathcal{F}$ is a P -point if it is an ultrafilter that is also a P -filter.

If $\mathcal{F}$ is a P -filter, then $\mathcal{F}$ is diagonalizable if and only if it is ω -diagonalizable if and only if it is ω -+-diagonalizable. On the other hand, these notions are distinct. Indeed the filter

$$\mathcal{F} = \mathfrak{F}\mathfrak{r} \otimes \mathfrak{F}\mathfrak{r} = \{X \subseteq \omega \times \omega : \{n : \{m : (n, m) \in X\} \text{ is cofinite}\} \text{ is cofinite}\}$$

is ω -diagonalizable, but not ω -+-diagonalizable; similarly, if $\mathcal{G}$ is any non-diagonalizable filter (any non-meager filter will do), then the filter

$$\mathcal{F} = \{X \subseteq \omega \times \omega : \text{for each } n, \{m : (n, m) \in X\} \in \mathcal{G}, \text{ and } \{n : \{m : (n, m) \in X\} \text{ is cofinite}\} \text{ is cofinite}\}$$

is ω -+-diagonalizable, but not diagonalizable.

If $\mathcal{F}$ is an ultrafilter, then $\mathcal{F}$ is a P -filter if and only if it is a weak P -filter if and only if it is a P^+ -filter; it is a Q -filter if and only if it is a weak Q -filter, and $\mathcal{F}$ is Ramsey if and only if it is weakly Ramsey if and only if it is +-Ramsey. Therefore these notions generalize the classical combinatorial properties of ultrafilters, but again, these notions can be seen to be different for filters in general.

The notions of ω -diagonalizability by $\mathcal{F}$ -(resp $\mathcal{F}^+$)-universal sets are generalizations of the regular ω -(resp. +)-diagonalizability. Observe that a filter $\mathcal{F}$ is ω -diagonalizable by $\mathfrak{F}\mathfrak{r}$ -universal sets if and only if diagonalizable by a single $\mathfrak{F}\mathfrak{r}$ -universal set if and only if it is meager, and ω -+-diagonalizability implies ω -diagonalizability by $\mathcal{F}$ -universal sets. This notion appears to be new.

Tree combinatorics is what most interest us in this paper as they naturally occur in terms of winning strategies; our main effort is then to express these combinatorics in terms of more familiar concepts. The following lemma shows the spirit of the paper.

Lemma 1.3. $\mathcal{F}$ is a non-meager P -filter if and only if every $\mathcal{F}$ -tree of finite sets has a branch whose union is in $\mathcal{F}$.

Proof: Assume first that every $\mathcal{F}$ -tree of finite sets has a branch whose union is in $\mathcal{F}$. Given a descending sequence $\langle A_n; n \in \omega \rangle \subseteq \mathcal{F}$, define an $\mathcal{F}$ -tree $\mathcal{T} \subseteq {}^{<\omega}([\omega]^{<\omega})$ such that $X_{\bar{s}} = A_n$ for each $\bar{s} \in \mathcal{T} \cap {}^n([\omega]^{<\omega})$. Any branch through $\mathcal{T}$ whose union is in $\mathcal{F}$ shows that $\mathcal{F}$ is a P -filter. To verify that $\mathcal{F}$ is non-meager, consider an increasing sequence of natural numbers $n_0 < n_1 < \dots$ and build again an $\mathcal{F}$ -tree of finite sets $\mathcal{T}$ as follows. Having already $\bar{s} \in \mathcal{T}$, choose k such that $\bar{s} \in {}^{<\omega}([n_k]^{<\omega})$ and let $X_{\bar{s}} = \omega \setminus n_{k+1}$. The union of any branch through $\mathcal{T}$ misses infinitely many intervals of the form $[n_k, n_{k+1})$ and therefore shows that $\mathcal{F}$ is also non-meager.

Now assume that $\mathcal{F}$ is a non-meager P -filter and let $\mathcal{T}$ be an $\mathcal{F}$ -tree of finite sets. Define $n_0 = 0$ and $A_0 = X_\emptyset$. More generally, given $A_0 \supseteq A_1 \supseteq \dots \supseteq A_k$ and $n_0 < n_1 < \dots < n_k$, let $n_{k+1} > n_k$ in A_k and put

$$A_{k+1} = \bigcap \{X_{\langle s_0, s_1, \dots, s_i \rangle} : i \leq k+1, s_j \subseteq A_j \cap [0, n_{k+1}]\},$$

Now as $\mathcal{F}$ is a P -filter, there is a $Y \in \mathcal{F}$ such that $Y \subseteq^* A_n$ for each n and we may as well assume by a reindexing that

$$Y \setminus n_{k+1} \subseteq A_k \text{ for each } k.$$

Now as $\mathcal{F}$ is also non-meager, we can find an infinite set $K = \{k_\ell : \ell \in \omega\}$ and we might as well assume that for each ℓ ,

$$Y \cap [n_{k_\ell}, n_{k_{\ell+1}}) = \emptyset.$$

Define $s_\ell = Y \cap [n_{k_\ell}, n_{k_{\ell+1}}) = Y \cap [n_{k_\ell+1}, n_{k_{\ell+1}})$ for each ℓ . We claim that $\langle s_k : k \in \omega \rangle$ is a branch through $\mathcal{T}$; indeed

$$\begin{aligned} s_\ell &= Y \cap [n_{k_\ell}, n_{k_{\ell+1}}) \\ &= Y \cap [n_{k_\ell+1}, n_{k_{\ell+1}}) \\ &\subseteq A_{k_\ell} \subseteq X_{\langle s_0, s_1, \dots, s_{\ell-1} \rangle}. \end{aligned}$$

But this concludes the proof as its union Y is in $\mathcal{F}$. $\square$

2. FILTER GAMES

We will be interested in infinite games of the form $\mathfrak{G}(\mathcal{X}, \mathcal{Y}, \mathcal{Z})$ where $\mathcal{X}$ will usually be a filter $\mathcal{F}$ or $\mathcal{F}^+$, $\mathcal{Y}$ will be either ω or $[\omega]^{<\omega}$, and $\mathcal{Z}$ will be either $\mathcal{F}$, $\mathcal{F}^+$, $\mathcal{F}^c$, the complement of $\mathcal{F}$ in $\wp(\omega)$ or else $(\mathcal{F}^+)^c = \mathcal{F}^*$, the dual ideal.

The game $\mathfrak{G}(\mathcal{X}, \mathcal{Y}, \mathcal{Z})$ is played by two players **I** and **II** as follows: at stage $k < \omega$, **I** chooses $X_k \in \mathcal{X}$, then **II** responds with either $n_k \in X_k$ in the case that $\mathcal{Y}$ is ω , or else responds with a nonempty $s_k \in [X_k]^{<\omega}$ in the case that $\mathcal{Y}$ is $[\omega]^{<\omega}$. At the end of the game, **II** is declared the winner if $\{n_k : k \in \omega\}$ (resp. $\bigcup_{k \in \omega} s_k$) $\in \mathcal{Z}$.

A few variations of these games have been considered in the literature, see for example in [2] and [11] for some special cases. In particular, the game $\mathfrak{G}(\mathfrak{Fr}, \omega, \mathcal{Z})$ is equivalent to the game in which at stage k player **I** chooses $m_k \in \omega$ and **II** responds with $n_k > m_k$, the outcome being that **II** wins the play if $\{n_k : k \in \omega\} \in \mathcal{Z}$ as before. Therefore we start with a result of [2].

Theorem 2.1. ([2]) Fix a filter $\mathcal{F}$ and consider the game $\mathfrak{G}(\mathfrak{Fr}, \omega, \mathcal{F})$. Then

1. **I** has a winning strategy if and only if $\mathcal{F}$ is not a Q -filter.
2. **II** has no winning strategy.

But this game generalizes in many ways and we have the following results.

Theorem 2.2. *Fix a filter $\mathcal{F}$ and consider the game $\mathfrak{G}(\mathfrak{F}\mathfrak{r}, \omega, \mathcal{F}^+)$. Then*

1. **I** has a winning strategy if and only if $\mathcal{F}$ is not a weak Q -filter .
2. **II** has a winning strategy if and only if $\mathcal{F}$ is ω -diagonalized.

A much less interesting game is the following:

Theorem 2.3. *Fix a filter $\mathcal{F}$ and consider the game $\mathfrak{G}(\mathfrak{F}\mathfrak{r}, \omega, \mathcal{F}^c)$. Then the game is determined and*

1. **II** always has a winning strategy.

Theorem 2.4. *Fix a filter $\mathcal{F}$ and consider the game $\mathfrak{G}(\mathfrak{F}\mathfrak{r}, \omega, \mathcal{F}^*)$. Then the game is determined and*

1. **I** has a winning strategy if and only if $\mathcal{F} = \mathfrak{F}\mathfrak{r}$ if and only if **II** has no winning strategy.

Because of the duality between $\mathfrak{F}\mathfrak{r}$ and $\mathfrak{F}\mathfrak{r}^+ = [\omega]^\omega$, we will get

Theorem 2.5. *Fix a filter $\mathcal{F}$, then the following games are dual of each other; that is a player has a winning strategy in one game if and only if the other player has a winning strategy in the other game.*

1. $\mathfrak{G}([\omega]^\omega, \omega, \mathcal{F})$ and $\mathfrak{G}(\mathfrak{F}\mathfrak{r}, \omega, \mathcal{F}^c)$.
2. $\mathfrak{G}([\omega]^\omega, \omega, \mathcal{F}^+)$ and $\mathfrak{G}(\mathfrak{F}\mathfrak{r}, \omega, \mathcal{F}^*)$.
3. $\mathfrak{G}([\omega]^\omega, \omega, \mathcal{F}^c)$ and $\mathfrak{G}(\mathfrak{F}\mathfrak{r}, \omega, \mathcal{F})$.
4. $\mathfrak{G}([\omega]^\omega, \omega, \mathcal{F}^*)$ and $\mathfrak{G}(\mathfrak{F}\mathfrak{r}, \omega, \mathcal{F}^+)$.

Now we consider the more interesting games where **I** plays members of the filter $\mathcal{F}$.

Theorem 2.6. *Fix a filter $\mathcal{F}$ and consider the game $\mathfrak{G}(\mathcal{F}, \omega, \mathcal{F})$. Then*

1. **I** has a winning strategy if and only if $\mathcal{F}$ is not a Ramsey filter.
2. **II** never has a winning strategy.

Theorem 2.7. *Fix a filter $\mathcal{F}$ and consider the game $\mathfrak{G}(\mathcal{F}, \omega, \mathcal{F}^+)$. Then*

1. **I** has a winning strategy if and only if $\mathcal{F}$ is not weakly Ramsey.
2. **II** has a winning strategy if and only if $\mathcal{F}$ is ω -+-diagonalizable.

Theorem 2.8. *Fix a filter $\mathcal{F}$ and consider the game $\mathfrak{G}(\mathcal{F}, \omega, \mathcal{F}^c)$. Then*

1. **I** never has a winning strategy.
2. **II** has a winning strategy if and only if $\mathcal{F}$ is not a Ramsey ultrafilter.

Theorem 2.9. *Fix a filter $\mathcal{F}$ and consider the game $\mathfrak{G}(\mathcal{F}, \omega, \mathcal{F}^*)$. Then*

1. **I** has a winning strategy if and only if $\mathcal{F}$ is countably generated.
2. **II** has a winning strategy if and only if $\mathcal{F}$ is not a +-Ramsey filter.

Again the duality of $\mathcal{F}$ and $\mathcal{F}^+$ will provide the following result.

Theorem 2.10. *Fix a filter $\mathcal{F}$, then the following games are dual of each other; that is a player has a winning strategy in one game if and only if the other player has a winning strategy in the other game.*

1. $\mathfrak{G}(\mathcal{F}^+, \omega, \mathcal{F})$ and $\mathfrak{G}(\mathcal{F}, \omega, \mathcal{F}^c)$.

2. $\mathfrak{G}(\mathcal{F}^+, \omega, \mathcal{F}^+)$ and $\mathfrak{G}(\mathcal{F}, \omega, \mathcal{F}^*)$.
3. $\mathfrak{G}(\mathcal{F}^+, \omega, \mathcal{F}^c)$ and $\mathfrak{G}(\mathcal{F}, \omega, \mathcal{F})$.
4. $\mathfrak{G}(\mathcal{F}^+, \omega, \mathcal{F}^*)$ and $\mathfrak{G}(\mathcal{F}, \omega, \mathcal{F}^+)$.

Before we turn to games in which player **II** chooses finite sets at each round, consider the following infinite game $\mathfrak{G}_1(\mathcal{F})$ defined in [2]: at stage k , player **I** chooses $m_k \in \omega$ and **II** responds with n_k . At the end, **II** is declared the winner if

1. $n_1 < n_2 < \dots < n_k < \dots$,
2. $m_k < n_k$ for infinitely many k , and
3. $\{n_k : k \in \omega\} \in \mathcal{F}$.

It is proved in [2] that **II** does not have any winning strategy in $\mathfrak{G}_1(\mathcal{F})$ and that **I** has a winning strategy if and only if $\mathcal{F}$ is meager. We have the following.

Theorem 2.11. *Fix a filter $\mathcal{F}$, the the games $\mathfrak{G}_1(\mathcal{F})$ and $\mathfrak{G}(\mathfrak{F}\mathfrak{r}, [\omega]^{<\omega}, \mathcal{F})$ are equivalent; that is a player has a winning strategy in one game if and only the same player has a winning strategy in the other game. Therefore, by [2], we have for either game,*

1. **I** has a winning strategy if and only if $\mathcal{F}$ is meager.
2. **II** never has a winning strategy.

Theorem 2.12. *Fix a filter $\mathcal{F}$, then the game $\mathfrak{G}(\mathfrak{F}\mathfrak{r}, [\omega]^{<\omega}, \mathcal{F}^+)$ is dual to the game $\mathfrak{G}(\mathfrak{F}\mathfrak{r}, [\omega]^{<\omega}, \mathcal{F})$. Therefore*

1. **I** never has a winning strategy.
2. **II** has a winning strategy if and only if $\mathcal{F}$ is meager.

As above, the following game is uninteresting.

Theorem 2.13. *Fix a filter $\mathcal{F}$, then the game $\mathfrak{G}(\mathfrak{F}\mathfrak{r}, [\omega]^{<\omega}, \mathcal{F}^c)$ is equivalent to $\mathfrak{G}(\mathfrak{F}\mathfrak{r}, \omega, \mathcal{F}^c)$. Therefore*

1. **II** always has a winning strategy.

Theorem 2.14. *Fix a filter $\mathcal{F}$, the the game $\mathfrak{G}(\mathfrak{F}\mathfrak{r}, [\omega]^{<\omega}, \mathcal{F}^*)$ is equivalent to $\mathfrak{G}(\mathfrak{F}\mathfrak{r}, \omega, \mathcal{F}^*)$. Therefore*

1. **I** has a winning strategy if and only if $\mathcal{F} = \mathfrak{F}\mathfrak{r}$ if and only if **II** has no winning strategy.

We turn to those games where **I** plays members of $\mathcal{F}$ and **II** responds with finite subsets.

Theorem 2.15. *Fix a filter $\mathcal{F}$ and consider the game $\mathfrak{G}(\mathcal{F}, [\omega]^{<\omega}, \mathcal{F})$. Then*

1. **I** has no winning strategy if and only if $\mathcal{F}$ is a non-meager P -filter.
2. **II** never has a winning strategy.

Theorem 2.16. *Fix a filter $\mathcal{F}$ and consider the game $\mathfrak{G}(\mathcal{F}, [\omega]^{<\omega}, \mathcal{F}^+)$. Then*

1. **I** has no winning strategy if and only if $\mathcal{F}$ is a weak P -filter.
2. **II** has a winning strategy if and only if $\mathcal{F}$ is ω -diagonalizable by $\mathcal{F}$ -universal sets.

Theorem 2.17. *Fix a filter $\mathcal{F}$, then the game $\mathfrak{G}(\mathcal{F}, [\omega]^{<\omega}, \mathcal{F}^c)$ is equivalent to the game $\mathfrak{G}(\mathcal{F}, \omega, \mathcal{F}^c)$. Therefore*

1. **I** never has a winning strategy.
2. **II** has a winning strategy if and only if $\mathcal{F}$ is not a Ramsey ultrafilter.

Theorem 2.18. *Fix a filter $\mathcal{F}$, then the game $\mathfrak{G}(\mathcal{F}, [\omega]^{<\omega}, \mathcal{F}^*)$ is equivalent to the game $\mathfrak{G}(\mathcal{F}, \omega, \mathcal{F}^*)$. Thus*

1. **I** has a winning strategy if and only if $\mathcal{F}$ is countably generated.
2. **II** has a winning strategy if and only if $\mathcal{F}$ is not a Ramsey ultrafilter.

Finally, we turn to games where **I** plays members of $\mathcal{F}^+$ while **II** replies with finite subsets. Note that we do not have here the same duality as when **II** responded with natural numbers.

Theorem 2.19. *Fix a filter $\mathcal{F}$ and consider the game $\mathfrak{G}(\mathcal{F}^+, [\omega]^{<\omega}, \mathcal{F})$. Then*

1. **I** has a winning strategy if and only if $\mathcal{F}$ is not a P -point.
2. **II** never has a winning strategy.

Theorem 2.20. *Fix a filter $\mathcal{F}$ and consider the game $\mathfrak{G}(\mathcal{F}^+, [\omega]^{<\omega}, \mathcal{F}^+)$.*

1. **I** has a winning strategy if and only if $\mathcal{F}$ is not a P^+ -filter.
2. **II** has a winning strategy if and only if $\mathcal{F}$ is ω -diagonalizable by $\mathcal{F}^+$ -universal sets.

Theorem 2.21. *Fix a filter $\mathcal{F}$, then the game $\mathfrak{G}(\mathcal{F}^+, [\omega]^{<\omega}, \mathcal{F}^c)$ is dual to the game $\mathfrak{G}(\mathcal{F}, \omega, \mathcal{F})$. Therefore*

1. **I** never has a winning strategy.
2. **II** has a winning strategy if and only if $\mathcal{F}$ is not a Ramsey filter.

Theorem 2.22. *Fix a filter $\mathcal{F}$, then the game $\mathfrak{G}(\mathcal{F}^+, [\omega]^{<\omega}, \mathcal{F}^*)$ is dual to the game $\mathfrak{G}(\mathcal{F}, \omega, \mathcal{F}^+)$. Therefore*

1. **I** has a winning strategy if and only if $\mathcal{F}$ is ω - $+$ -diagonalizable.
2. **II** has a winning strategy if and only if $\mathcal{F}$ is not weakly Ramsey.

3. PROOFS

In this section we verify the results of section 2. We start with two general results.

Lemma 3.1. *If a family $\mathcal{Z} \subseteq \wp(\omega)$ is closed under supersets, then the two games $\mathfrak{G}(\mathcal{X}, \omega, \mathcal{Z}^c)$ and $\mathfrak{G}(\mathcal{X}, [\omega]^{<\omega}, \mathcal{Z}^c)$ are equivalent.*

Proof: Since player **II** is trying to get out of $\mathcal{Z}$ which is assumed to be closed under supersets (therefore $\mathcal{Z}^c$ is closed under subsets), the best strategy for **II** is to play finite sets as small as possible, namely singleton since a legal move must be nonempty. $\square$

The next Lemma regards the duality mentioned in the introduction and is taken from the work of Aczel [1] and Blass (see [3], Theorem 1). We include a hint of the proof for completeness.

Theorem 3.2. *([1],[3]) For a given filter $\mathcal{F}$, the game $\mathfrak{G}(\mathcal{F}, \omega, \mathcal{Z})$ and the game $\mathfrak{G}(\mathcal{F}^+, \omega, \mathcal{Z}^c)$ are dual; that is a player has a winning strategy in one of these games if and only if the other player has a winning strategy in the other game.*

Proof: Suppose **II** has a winning strategy $\$$ in the game $\mathfrak{G}(\mathcal{F}, \omega, \mathcal{Z})$, we define a strategy $\$$ for **I** in the game $\mathfrak{G}(\mathcal{F}^+, \omega, \mathcal{Z}^c)$ as follows:

I starts with $\$(\emptyset) = \{ \$(X) : X \in \mathcal{F} \} \in \mathcal{F}^+$. When **II** responds with n_0 , **I** remembers one set $X_0 \in \mathcal{F}$ such that $\$(X_0) = n_0$.

At stage k , **I** has remembered k sets $X_0, X_1, \dots, X_{k-1}$ from $\mathcal{F}$ while **II** responded with $\langle n_0, n_1, \dots, n_{k-1} \rangle$. **I** then plays

$$\$(\langle n_0, n_1, \dots, n_{k-1} \rangle) = \{ \$(\langle X_0, X_1, \dots, X_{k-1}, X \rangle) : X \in \mathcal{F} \} \in \mathcal{F}^+;$$

II responds with n_k and **I** remembers one set $X_k \in \mathcal{F}$ such that

$$\$(\langle X_0, X_1, \dots, X_{k-1}, X_k \rangle) = n_k.$$

Thus a play in the new game corresponds to a play in the former game and thus the outcome $\{n_k : k \in \omega\} \in \mathcal{Z}$ and **I**'s strategy is a winning strategy in $\mathfrak{G}(\mathcal{F}^+, \omega, \mathcal{Z}^c)$.

The other cases are quite similar and left to the reader. $\square$

Now we are ready to attack the proofs of section 2.

PROOF OF THEOREM 2.2: We first deal with player **I**. So suppose that $\mathcal{F}$ is not a weak Q-filter and therefore there is a partition of ω into finite sets $\langle s_k : k \in \omega \rangle$ such that no $X \in \mathcal{F}^+$ meet each s_k in at most one point. Then **I**'s strategy at stage k , after **II** has played $\langle n_0, n_1, \dots, n_{k-1} \rangle$, is to respond with $\bigcup \{s_i : s_i \cap \{n_0, n_1, \dots, n_{k-1}\} = \emptyset\}$.

Now suppose that $\mathcal{F}$ is a weak Q-filter and we show that any strategy $\$$ for **I** is not a winning strategy. Define a sequence of integers $\langle \pi_k : k \in \omega \rangle$ such that $[\pi_0, \infty) \subseteq \$(\emptyset)$ and more generally

$$[\pi_{k+1}, \infty) \subseteq \bigcap \{ \$(\langle n_0, n_1, \dots, n_i \rangle) : n_0 < n_1 < \dots < n_i < \pi_k \}.$$

By assumption there is an $X \in \mathcal{F}^+$ which meets each interval $[\pi_k, \pi_{k+1})$ in at most one point. But $X = X_0 \cup X_1$ where $X_i = X \cap \bigcup_k [\pi_{2k+i}, \pi_{2k+i+1})$ and therefore $X_i \in \mathcal{F}^+$ for some i . Write X_i in increasing order as $\langle n_k : k \in \omega \rangle$; but then X_i is the outcome of a legal play won by **II**, and thus $\$$ was not a winning strategy for **I**.

Now we deal with player **II**. Suppose that $\mathcal{F}$ is ω -diagonalized by $\langle X_n : n \in \omega \rangle \subseteq [\omega]^\omega$. Fix a surjective map $\sigma : \omega \rightarrow \omega$ such that the preimage of each n is infinite. Here is **II**'s strategy: at stage k , after **I** played $Y_k \in \mathfrak{F}\mathfrak{r}$, **II** responds with an element of $Y_k \cap X_{\sigma(k)} \setminus k$. At the end of the play, **II**'s outcome is a set with infinite intersection with each X_n and therefore belongs to $\mathcal{F}^+$, thus this is a winning strategy for **II**.

Now let $\$$ be a winning strategy for **II** in the game, we show that $\mathcal{F}$ is ω -diagonalizable. We claim that

$$(\forall Y \in \mathcal{F})(\exists n = n(Y))(\exists s = s(Y) \in {}^n \omega)(\forall^\infty t \in {}^{n+1} \omega) s < t \implies \$(t) \in Y.$$

Indeed otherwise one quickly produces a winning play for **I**. But then the collection $\{ \$(s^n); n \in \omega \}; s \in {}^{<\omega} \omega \}$ ω -diagonalize $\mathcal{F}$. This completes the proof. $\square$

PROOF OF THEOREM 2.3: This is trivial; **II** chooses an infinite $X \notin \mathcal{F}$ and plays continually members of X . $\square$

PROOF OF THEOREM 2.4: If $\mathcal{F} = \mathfrak{F}\mathfrak{r}$, then $\mathcal{F}^+ = [\omega]^\omega$ and **I**'s strategy is to ensure that **II**'s outcome is infinite, and of course **II** has no winning strategy.

If however $\mathcal{F} \neq \mathfrak{F}\mathfrak{r}$, **II** chooses an infinite $X \notin \mathcal{F}^+$ and continually plays members of X ; this constitutes a winning strategy for **II**. $\square$

PROOF OF THEOREM 2.5: This follows from Theorem 3.2 as $\mathfrak{F}\mathfrak{r}^+ = [\omega]^\omega$. $\square$

PROOF OF THEOREM 2.6: If **II** had a winning strategy in this game, it would also be a winning strategy for $\mathfrak{G}(\mathfrak{F}\mathfrak{r}, \omega, \mathcal{F})$, which is impossible.

Now for player **I**. If $\mathcal{F}$ is not a Q-filter, then **I** fixes a partition of ω into finite sets $\langle s_k : k \in \omega \rangle$ and plays to ensure that **II**'s outcome meets each s_k in at most one point; therefore **I** wins. If on the other hand $\mathcal{F}$ is not a P-filter, **I** then fixes a sequence $\langle X_n : n \in \omega \rangle \subseteq \mathcal{F}$ such that no $X \in \mathcal{F}$ is almost included in each X_n . It suffices for **I** to play $\cap_{i < k} X_i$ at stage k to produce a winning strategy.

This leaves us with the more interesting situation in which $\mathcal{F}$ is a Q-filter P-filter and we must show that any strategy $\$$ for **I** cannot be a winning one.

Fixing such a strategy $\$,$ **II** first chooses $Y \in \mathcal{F}$ such that $Y \subseteq^* \$(s)$ for all $s \in {}^{<\omega}\omega$. Now **II** defines a sequence $\langle X_n : n \in \omega \rangle \subseteq \mathcal{F}$ and a sequence $\langle n_k : k \in \omega \rangle$ as follows: $X_0 = \$(\emptyset)$ and n_0 is such that $Y \setminus n_0 \subseteq X_0$. Now given $n_0 < n_1 < \dots < n_k$, put

$$X_{k+1} = \bigcap \{ \$(\langle m_0, m_1, \dots, m_i \rangle) : i \leq k, m_j \in [n_j, n_k] \cap Y \}.$$

Then **II** chooses n_{k+1} such that $Y \setminus n_{k+1} \subseteq X_{k+1}$. Now because $\mathcal{F}$ is a Q-filter, **II** knows very well that there is a set in $\mathcal{F}$ missing infinitely many intervals, say $[n_{k_n}, n_{k_n+1})$, for an infinite set $\{k_n : n \in \omega\}$, and therefore by selectivity again **II** can find a set $Y' = \{y_n : n \in \omega\} \subseteq Y$ in $\mathcal{F}$ such that Y' misses all these intervals $[n_{k_n}, n_{k_n+1})$ and further intersects each other interval $[n_{k_n}, n_{k_n+1})$ in at most one point.

But we claim now that Y' is a legal play in the game! Indeed, for each k , say $y_k \in [n_{k_n}, n_{k_n+1}) \cap Y' = [n_{k_n+1}, n_{k_n+1}) \cap Y'$, then $y_k \in X_{k_n+1} \subseteq \$(\langle y_0, y_1, \dots, y_{k-1} \rangle)$. Thus **I**'s strategy $\$$ was definitely not a winning one. $\square$

PROOF OF THEOREM 2.7: We first look at player **I**. If $\mathcal{F}$ is not weakly Ramsey, there is an $\mathcal{F}$ -tree $\mathcal{T}$ such that no branches belong to $\mathcal{F}^+$; this gives a winning strategy for **I** by playing along the tree.

Now suppose that $\mathcal{F}$ is weakly Ramsey and that $\$$ is a strategy for **I**, we will produce a winning play for **II** showing that **I** cannot have a winning strategy. Let $X_0 = \$(\emptyset)$ and choose $n_0 \in X_0$. Having produced $n_0 < n_1 < \dots < n_k$, let

$$X_{k+1} = \bigcap \{ \$(y_0, y_1, \dots, y_i) : i \leq k, y_0 < y_1 < \dots < y_k \leq n_k \} \setminus n_k.$$

$X_{k+1} \in \mathcal{F}$ and choose $n_{k+1} \in X_{k+1}$. Now define a $\mathcal{F}$ -tree $\mathcal{T}$ inductively by letting $X_\emptyset = X_0$, and given $s = \langle y_0, y_1, \dots, y_i \rangle \in \mathcal{T}$, say $n_{k-1} \leq y_i < n_k$, then let $X_s = X_{k+1}$. This $\mathcal{F}$ -tree $\mathcal{T}$ must contain a branch $\{y_k : k \in \omega\} \in \mathcal{F}^+$ by assumption, but this clearly is a legal play of the game in which **II** wins.

Now we deal with player **II**. Suppose that $\mathcal{F}$ is ω -+-diagonalized by $\langle X_n : n \in \omega \rangle \in \mathcal{F}^+$. Fix a surjective map $\sigma : \omega \rightarrow \omega$ such that the preimage of each n is infinite. Here is **II**'s strategy: at stage k , after **I** played $Y_k \in \mathcal{F}$, **II** responds with an element of $Y_k \cap X_{\sigma(k)} \setminus k$. Notice the importance of having each $X_n \in \mathcal{F}^+$. At the end of the play, **II**'s outcome is a set with infinite intersection with each X_n and therefore belongs to $\mathcal{F}^+$, thus it is a winning strategy for **II**.

Now let $\$$ be a winning strategy for **II** in the game, we show that $\mathcal{F}$ is ω -+-diagonalizable. We first define an $\mathcal{F}^+$ -tree as follows. Let $X_\emptyset = \{ \$(X) : X \in \mathcal{F} \} \in$

$\mathcal{F}^+$, and for each $n \in X_\emptyset$, select an $X_\emptyset^n \in \mathcal{F}$ such that $\$(X_\emptyset^n) = n$. More generally, given $X_{\vec{s}}^n \in \mathcal{F}$, for $n \in X_{\vec{s}} \in \mathcal{F}^+$, say $\vec{s} = \langle s_0, s_1, \dots, s_i \rangle$, let

$$X_{\vec{s}} \wedge n = \{ \$(X_\emptyset^{s_0}, X_{\langle s_0 \rangle}^{s_1}, X_{\langle s_0, s_1 \rangle}^{s_2}, \dots, X_{\langle s_0, s_1, \dots, s_{i-1} \rangle}^{s_i}, X) : X \in \mathcal{F} \} \in \mathcal{F}^+,$$

and for each $k \in X_{\vec{s}} \wedge n$, choose $X_{\vec{s}}^k \wedge n \in \mathcal{F}$ such that

$$\$(X_\emptyset^{s_0}, X_{\langle s_0 \rangle}^{s_1}, X_{\langle s_0, s_1 \rangle}^{s_2}, \dots, X_{\langle s_0, s_1, \dots, s_{i-1} \rangle}^{s_i}, X_{\vec{s}}^k \wedge n) = k.$$

Therefore we obtain a $\mathcal{F}^+$ -tree $\mathcal{T}$ each of whose branches is a legal play of the game, and therefore belongs to $\mathcal{F}^+$ as $\$$ was a winning strategy for **II** and this means that the sets $\{X_{\vec{s}} : \vec{s} \in \mathcal{T}\} \subseteq \mathcal{F}^+$ must ω -+-diagonalize $\mathcal{F}$. $\square$

PROOF OF THEOREM 2.8: As in $\mathfrak{G}(\mathfrak{F}\mathfrak{r}, \omega, \mathcal{F}^c)$, **I** has no winning strategy.

As for **II**, let us consider the case when $\mathcal{F}$ is not a Ramsey ultrafilter first. There are three possibilities which **II** figures out. If $\mathcal{F}$ is not an ultrafilter, **II** chooses $X \in \mathcal{F}^+ \setminus \mathcal{F}$ and constantly plays members of X , and therefore wins the game. Otherwise **II** checks whether $\mathcal{F}$ is a Q-filter and if not chooses a partition $\langle s_k : k \in \omega \rangle$ of ω into finite sets for which $\mathcal{F}$ contains no selector. But then **II**'s strategy is to play a selector for the partition, therefore winning again. Finally **II** realizes that it must be that $\mathcal{F}$ is not a P-filter and thus selects $\langle X_n : n \in \omega \rangle \subseteq \mathcal{F}$ with no $Y \in \mathcal{F}$ almost included in each X_n . Then **II**'s strategy at the k^{th} move is to play a member of $\cap_{i < k} X_i$ and again has a winning strategy.

Now we suppose that $\mathcal{F}$ is a Ramsey ultrafilter and must show that **II** cannot have a winning strategy. But as $\mathcal{F}$ is an ultrafilter, $\mathcal{F} = \mathcal{F}^+$ and by the duality theorem 3.2, the game $\mathfrak{G}(\mathcal{F}, \omega, \mathcal{F}^c)$ is dual to the game $\mathfrak{G}(\mathcal{F}, \omega, \mathcal{F})$, and as $\mathcal{F}$ is a Q-filter P-filter, **I** has no winning strategy in $\mathfrak{G}(\mathcal{F}, \omega, \mathcal{F})$ by theorem 2.6 and therefore **II** has no winning strategy in $\mathfrak{G}(\mathcal{F}, \omega, \mathcal{F}^c)$. $\square$

PROOF OF THEOREM 2.9: As far as player **I** is concerned, if $\mathcal{F}$ is generated by countably many sets $\langle X_n : n \in \omega \rangle$, then it suffices for **I** to play $\cap_{i < k} X_i \setminus k$ at stage k , and thus the outcome of the play is a set $Y \subseteq^* X_n$ for each n , definitely in $\mathcal{F}^+$ and **I** wins.

If on the other hand **I** has a winning strategy $\$$ in the game, then the filter $\mathcal{F}$ must be generated by $\langle \$(s) : s \in {}^{<\omega}\omega \rangle$, as can easily be verified.

Now for player **II**. If $\mathcal{F}$ is not +-Ramsey, then there is an $\mathcal{F}^+$ -tree $\mathcal{T}$ none of whose branches belong to $\mathcal{F}^+$; therefore **II**'s strategy is to play along the tree $\mathcal{T}$. Suppose finally that $\mathcal{F}$ is +-Ramsey and we show that any strategy $\$$ for **II** is not a winning strategy.

We first define a $\mathcal{F}^+$ -tree as follows. Let $X_\emptyset = \{ \$(X) : X \in \mathcal{F} \} \in \mathcal{F}^+$, and for each $n \in X_\emptyset$, select an $X_\emptyset^n \in \mathcal{F}$ such that $\$(X_\emptyset^n) = n$. More generally, given $X_{\vec{s}}^n \in \mathcal{F}$, for $n \in X_{\vec{s}} \in \mathcal{F}^+$, say $\vec{s} = \langle s_0, s_1, \dots, s_i \rangle$, let

$$X_{\vec{s}} \wedge n = \{ \$(X_\emptyset^{s_0}, X_{\langle s_0 \rangle}^{s_1}, X_{\langle s_0, s_1 \rangle}^{s_2}, \dots, X_{\langle s_0, s_1, \dots, s_{i-1} \rangle}^{s_i}, X) : X \in \mathcal{F} \} \in \mathcal{F}^+,$$

and for each $k \in X_{\vec{s}} \wedge n$, choose $X_{\vec{s}}^k \wedge n \in \mathcal{F}$ such that

$$\$(X_\emptyset^{s_0}, X_{\langle s_0 \rangle}^{s_1}, X_{\langle s_0, s_1 \rangle}^{s_2}, \dots, X_{\langle s_0, s_1, \dots, s_{i-1} \rangle}^{s_i}, X_{\vec{s}}^k \wedge n) = k.$$

Therefore we obtain a $\mathcal{F}^+$ -tree $\mathcal{T}$ all of whose branches are a legal play of the game, but there is such a branch in $\mathcal{F}^+$ as $\mathcal{F}$ is +-Ramsey and therefore **II**'s strategy is not a winning strategy. $\square$

PROOF OF THEOREM 2.10: Follows immediately from Theorem 3.2. $\square$

PROOF OF THEOREM 2.11: We must show that the games $\mathfrak{G}_1(\mathcal{F})$ is equivalent to the game $\mathfrak{G}(\mathfrak{F}\mathfrak{r}, [\omega]^{<\omega}, \mathcal{F})$. As **I** is trying to produce an outcome out of $\mathcal{F}$ in the game $\mathfrak{G}(\mathfrak{F}\mathfrak{r}, [\omega]^{<\omega}, \mathcal{F})$, **I** plays without loss of generality cofinite sets of the form $[n, \infty)$ which we identify with n to simplify notation.

Suppose first that **I** has a winning strategy $\$$ in the game $\mathfrak{G}_1(\mathcal{F})$, we define a strategy $\bar{\$}$ for **I** in the game $\mathfrak{G}(\mathfrak{F}\mathfrak{r}, [\omega]^{<\omega}, \mathcal{F})$ by

$$\bar{\$}(s_0, s_1, \dots, s_i) = \$\left(\bigcup_{j \leq i} s_j\right),$$

where $(\bigcup_{j \leq i} s_j)$ is considered as an element of $[\omega]^{<\omega}$. This gives a winning strategy for **I** in $\mathfrak{G}(\mathfrak{F}\mathfrak{r}, [\omega]^{<\omega}, \mathcal{F})$.

Now let $\$$ be a winning strategy for **I** in $\mathfrak{G}(\mathfrak{F}\mathfrak{r}, [\omega]^{<\omega}, \mathcal{F})$, we define a strategy $\bar{\$}$ for **I** in $\mathfrak{G}_1(\mathcal{F})$ as follows:

at stage k , after **I** played $m_0, m_1, \dots, m_k$ and **II** responded with $n_0, n_1, \dots, n_k$, **I** puts $s_i = (m_{i-1}, m_i] \cap \{n_0, n_1, \dots, n_k\}$ for $i \leq k$ and $s_{k+1} = [m_k, \infty) \cap \{n_0, n_1, \dots, n_k\}$ and replies with $\$(s_0, s_1, \dots, s_{k+1})$. This must be a winning strategy for **I**. Indeed let $m_0 = \$(\emptyset)$ be the first move of **I**. Now given m_k , there must be a stage at which **II** responded with an integer bigger than **I**'s play, as otherwise **I** wins the play trivially, so call this last move of **I** by m_{k+1} . Then if s_k denotes **II**'s moves in the interval $[m_k, m_{k+1})$, then we get a legal play $\langle m_0, s_0, m_1, s_1, \dots \rangle$ in the game $\mathfrak{G}(\mathfrak{F}\mathfrak{r}, [\omega]^{<\omega}, \mathcal{F})$ which **I** wins by assumptions; but as the outcome is the same we have shown that $\bar{\$}$ is a winning strategy for **I** in the game $\mathfrak{G}_1(\mathcal{F})$.

Now we deal with player **II**. If $\$$ is a winning strategy for **II** in the game $\mathfrak{G}(\mathfrak{F}\mathfrak{r}, [\omega]^{<\omega}, \mathcal{F})$, then **II** behaves as follows in the game $\mathfrak{G}_1(\mathcal{F})$: at stage k , **II** imagines that **I** has played $\langle m_0, m_1, \dots, m_k \rangle$ in the game $\mathfrak{G}(\mathfrak{F}\mathfrak{r}, [\omega]^{<\omega}, \mathcal{F})$ and plays one by one the elements of $\$(\langle m_0, m_1, \dots, m_k \rangle)$ without noticing **I**'s moves until done, and then remembers **I**'s last move, m_{k+1} . At the end, the outcome is one of the game $\mathfrak{G}(\mathfrak{F}\mathfrak{r}, [\omega]^{<\omega}, \mathcal{F})$ and thus **II** wins.

Finally suppose that $\$$ is a winning strategy for **II** in the game $\mathfrak{G}_1(\mathcal{F})$.

Claim 3.3. [2] *For each $\sigma \in [\omega]^{<\omega}$, there is $\tau \in [\omega]^{<\omega}$ and $n \in \omega$, such that $\$(\sigma \wedge \tau \wedge m) > m$ for each $m > n$.*

If the claim were false, then it is left to the reader to verify that **I** could easily win a play.

So here is what **II** does in the game $\mathfrak{G}(\mathfrak{F}\mathfrak{r}, [\omega]^{<\omega}, \mathcal{F})$: at stage k , after **I** has played $\sigma = \langle m_0, m_1, \dots, m_k \rangle$, **II** chooses the corresponding $\tau = \langle t_0, t_1, \dots, t_\ell \rangle$ and n_k from the lemma and replies with $s_k = \$\sigma \wedge (\tau \upharpoonright i)$ for $i \leq \ell$; when **I** responds with m_{k+1} , **II** just imagines that **I** actually responded with some $\bar{m}_{k+1} > n_k$. This gives a simulated play from $\mathfrak{G}_1(\mathcal{F})$ and therefore **II** wins. $\square$

PROOF OF THEOREM 2.12: The fact that this game is dual to $\mathfrak{G}(\mathfrak{F}\mathfrak{r}, [\omega]^{<\omega}, \mathcal{F})$ is purely accidental I believe.

By playing two games simultaneously, **II** can produce two outcomes whose union is cofinite, therefore one of them must be in $\mathcal{F}^+$ and thus **I** has no winning strategy. More precisely, let $m_0 = \bar{m}_0$ be **I**'s first move as we again identify a cofinite set $[m, \infty)$ with m . **II** replies with $\{m_0\}$, and then **I** responds with m_1 in the first game. Now in the second game **II** replies with $[m_0, m_1]$ and waits for **I**'s response $\bar{m}_1$;

then **II** comes back to the first game and replies with $[m_1, \overline{m}_1]$. Continuing this way, **II** produces the outcome $A = \bigcup_i [m_i, \overline{m}_i]$ in the first game, and $B = \bigcup_i [\overline{m}_i, m_{i+1}]$ in the second. One of these sets is in $\mathcal{F}^+$ and therefore **I** lost one of the games.

Now for player **II**. If $\mathcal{F}$ is meager, then **II** has definitely an easy time winning the game; indeed there must be a sequence $\pi_0 < \pi_1 < \dots$ such that each member of $\mathcal{F}$ meets all but finitely many of the intervals $[\pi_k, \pi_{k+1})$. Therefore at stage ℓ , **II** plays one of the intervals $[\pi_k, \pi_{k+1})$ with $k > \ell$.

So we must show this is the only way that **II** can have a winning strategy. So fix such a winning strategy $\$$ for **II**. Define a sequence of integers by $\pi_0 = 1$, and given π_k , choose

$$\pi_{k+1} = \max\{\$(m_0, m_1, \dots, m_i) : i \leq k \text{ and } m_j \leq \pi_k\} + 1.$$

then each member of $\mathcal{F}$ must meet all but finitely many of the intervals $[\pi_k, \pi_{k+1})$. Otherwise, say $Y \in \mathcal{F}$ misses the intervals $[\pi_{k_\ell}, \pi_{k_\ell+1})$ for $\ell \in \omega$, then **I** wins by playing exactly these k_ℓ . $\square$

PROOF OF THEOREM 2.13: Follows immediately by Lemma 3.1 and Theorem 2.3. $\square$

PROOF OF THEOREM 2.14: Follows immediately by Lemma 3.1 and Theorem 2.4. $\square$

PROOF OF THEOREM 2.15: If **II** had a winning strategy in this game, it would be a winning strategy in the game $\mathfrak{G}(\mathfrak{F}\mathfrak{r}, [\omega]^{<\omega}, \mathcal{F})$, which is impossible.

If $\mathcal{F}$ is meager, then **I** uses the same strategy as for $\mathfrak{G}(\mathfrak{F}\mathfrak{r}, [\omega]^{<\omega}, \mathcal{F})$; if on the other hand $\mathcal{F}$ is not a P-filter, then **I** chooses a witness $\langle X_n : n \in \omega \rangle$ and plays $\bigcap_{i < k} X_i$ at stage k which provides a winning strategy.

Now, as a strategy for player **I** is nothing else but an $\mathcal{F}$ -tree of finite sets, it should be rather clear that **I** has no winning strategy if and only if any $\mathcal{F}$ -tree has a branch in $\mathcal{F}$, that is if and only if $\mathcal{F}$ is a non-meager P-filter by Lemma 1.3. $\square$

PROOF OF THEOREM 2.16: As far as player **I** is concerned, the proof is entirely similar to that of Theorem 2.7.

So we consider the situation for **II**. First fix $\mathcal{F}$ -universal sets $\langle X_n : n \in \omega \rangle$ ω -diagonalizing $\mathcal{F}$. Player **II** fixes a surjection $\sigma : \omega \rightarrow \omega$ such that the preimage of every n is infinite and at stage k , after **I** produced a set $Y_k \in \mathcal{F}$, **II** responds with $s_k \in X_{\pi(k)} \cap [Y_k \setminus k]^{<\omega}$. At the end of the play, **II** has produced $S = \bigcup_k s_k$ which contains infinitely many members of each X_n , and is therefore in $\mathcal{F}^+$.

Now let $\$$ be a winning strategy for **II** and we define a tree $\mathcal{T} \subseteq {}^{<\omega}([\omega]^{<\omega})$ such that the successors of each node $\overline{s} \in \mathcal{T}$ form an $\mathcal{F}$ -universal set and the collection $\langle X_{\overline{s}} : \overline{s} \in \mathcal{T} \rangle$ ω -diagonalizes $\mathcal{F}$.

Let $X_\emptyset = \{\$(X) : X \in \mathcal{F}\}$, an $\mathcal{F}$ -universal set, and for each $s \in X_\emptyset$, choose $X_\emptyset^s \in \mathcal{F}$ such that $\$(X_\emptyset^s) = s$. In general, given $X_{\overline{s}}^t \in \mathcal{F}$, for $\overline{s} = \langle s_0, s_1, \dots, s_i \rangle \in {}^{<\omega}([\omega]^{<\omega})$, define

$$X_{\overline{s}}^t \wedge t = \{\$(X_\emptyset^{s_0}, X_{s_0}^{s_1}, \dots, X_{\overline{s}}^t, X) : X \in \mathcal{F}\}$$

a $\mathcal{F}$ -universal set, and for each $u \in X_{\overline{s}}^t \wedge t$, choose $X_{\overline{s}}^u \wedge t \in \mathcal{F}$ such that

$$\$(X_\emptyset^{s_0}, X_{s_0}^{s_1}, \dots, X_{\overline{s}}^t, X_{\overline{s}}^u \wedge t) = u.$$

Now every branch of $\mathcal{T}$ constitutes an outcome of a play of the game, and therefore the sets $(\mathcal{F}\text{-universal}) X_{\bar{s}}$ for $\bar{s} \in \mathcal{T}$ must ω -diagonalize $\mathcal{F}$. $\square$

PROOF OF THEOREM 2.17: Follows immediately from Lemma 3.1 and Theorem 2.8. $\square$

PROOF OF THEOREM 2.18: Follows immediately from Lemma 3.1 and Theorem 2.9. $\square$

PROOF OF THEOREM 2.19: Player **II** definitely has no hope for a winning strategy; if there is $X \in \mathcal{F}^+ \setminus \mathcal{F}$, then **II** is doomed, and otherwise $\mathcal{F} = \mathcal{F}^+$ and the result follows by Theorem 2.15.

So player **I** has a winning strategy if $\mathcal{F}^+ \neq \mathcal{F}$, and otherwise, the result follows again from Theorem 2.15. $\square$

PROOF OF THEOREM 2.20: The proof is entirely similar to that of Theorem 2.16 and is left to the reader.

PROOF OF THEOREM 2.21: Follows immediately from Lemma 3.1, Theorem 3.2 and Theorem 2.6. $\square$

PROOF OF THEOREM 2.22: Follows immediately from Lemma 3.1, Theorem 3.2 and Theorem 2.7. $\square$

4. CONCLUSION

The combinatorial properties of a filter being meager, a P-filter or a Q-filter have probably been the most popular; another related property which has been around for some times is that of a filter $\mathcal{F}$ being ‘rapid’, i.e. for any partition of ω into finite sets $\langle s_k : k \in \omega \rangle$, $\mathcal{F}$ contains a set X such that $X \cap s_k$ has size at most k for each k . If we modify the games above so that at stage k player **II** responds with a finite set of size at most k (or bounded by a fixed “unbounded” function), then the characterizations of the winning strategies for either player will involve modifications of this property of being rapid, such as weakly rapid and so on. The details are left to the interested reader.

However, we have little information on the following interesting variation of the games:

Problem: Characterize winning strategies for the games in which player **II** responds with members of $\mathcal{F}^*$.

REFERENCES

1. P. Aczel, *Quantifiers, Games, and Inductive Definitions*, in Proc. Third Scandinavian Logic Symposium, ed. S. Kanger, North Holland, 1975, 1-14.
2. T. Bartoszyński and M. Scheepers, *Filter and Games*, to appear, 1993.
3. A. Blass, *Selective Ultrafilters and Homogeneity*, Annals of Pure and Applied Logic **38** (1988), 215-255.
4. D. Booth, *Ultrafilters on a Countable Set*, Annals Math. Logic **2** (1970), 1-24.
5. F. Galvin, Unpublished manuscript on ultrafilter games, $\approx$ 1980.
6. S. Grigorieff, *Combinatorics on Ideals and Forcing*, Annals Math. Logic **3** (1971), 363-394.
7. W. Just, A.R.D. Mathias, K. Prikry and P. Simon, *On the existence of large P-ideals*, to appear.
8. K. Kunen, *Some Points in βN* , Math. Proc. Cambridge Phil. Soc. **80** (1976), 385-398.
9. C. Laflamme, *Zapping Small Filters*, Proc. Amer. Math. Soc. **114** (1992), 535-544.

10. ———, *Strong Meager Properties for Filter*, 1993.
11. S. Shelah, *Proper Forcing*, Lecture Notes in Mathematics, Springer-Verlag, Berlin 1982.
12. M. Talagrand, *Compacts de fonctions mesurables et filtres non mesurables*, Studia Math. **67** (1980), 13-43.

DEPARTMENT OF MATHEMATICS AND STATISTICS, UNIVERSITY OF CALGARY, CALGARY, ALBERTA, CANADA T2N 1N4

E-mail address: laflamme@acs.ucalgary.ca